

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8/18(K8.8 + DC10), 1mg/mL, Clone: [K8.8 DC10], Mouse, Monoclonal BOT-BNUM0691-50

Artikelname	Cytokeratin 8/18(K8.8 + DC10), 1mg/mL, Clone: [K8.8 DC10], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0691-50
Hersteller Artikelnummer	BNUM0691-50
Alternativnummer	BOT-BNUM0691-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Keratin preparation from a human carcinoma (K8.8), PMC-42 human breast carcinoma cells (DC10)
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb cocktail recognizes all simple epithelia including glandular epithelium, for exa...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[K8.8 DC10]

Molekulargewicht	52.5 kDa (CK8), 45 kDa (CK18)
UniProt	P05787
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user