

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD57 / B3GAT1(NK/804), 1mg/mL, Clone: [NK/804], Mouse, Monoclonal BOT-BNUM0804-50

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD57 / B3GAT1(NK/804), 1mg/mL, Clone: [NK/804], Mouse, Monoclonal                                                                                                                                                                                             |
| Artikelnummer            | BOT-BNUM0804-50                                                                                                                                                                                                                                               |
| Hersteller Artikelnummer | BNUM0804-50                                                                                                                                                                                                                                                   |
| Alternativnummer         | BOT-BNUM0804-50-50UL                                                                                                                                                                                                                                          |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human B3GAT1 protein                                                                                                                                                                                                                              |
| Produktbeschreibung      | Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including c... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 1 mg/mL                                                                                                                                                                                                                                                       |
| Klon-Bezeichnung         | [NK/804]                                                                                                                                                                                                                                                      |
| Molekulargewicht         | ~110 kDa                                                                                                                                                                                                                                                      |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt                | <a href="#">Q9P2W7</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Puffer                 | PBS, no BSA, no azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |