

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytochrome C (Mitochondrial Marker)(7H8.2C12 + CYCS/1010), 1mg/mL, Clone: [7H8.2C12 CYCS/1010], Mouse, Monoclonal BOT-BNUM1265-50

Artikelname	Cytochrome C (Mitochondrial Marker)(7H8.2C12 + CYCS/1010), 1mg/mL, Clone: [7H8.2C12 CYCS/1010], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1265-50
Hersteller Artikelnummer	BNUM1265-50
Alternativnummer	BOT-BNUM1265-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC, WB
Spezies Reaktivität	Human, Rat
Immunogen	Synthetic peptides corresponding to amino acid 1-80, 81-104 and 66-104 of pigeon cytochrome c (7H8.2C12), Recombinant full-length human CYCS protein (CYCS/1010)
Produktbeschreibung	Cytochrome C is a well-characterized mobile electron transport protein that is essential to energy conversion in all aerobic organisms. In mammalian cells, this highly conserved protein is normally localized to the mitochondrial inter-membrane space....
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[7H8.2C12 CYCS/1010]

Molekulargewicht	15 kDa
UniProt	P99999
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.25-0.5 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Western Blot 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user