

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tyrosinase (Melanoma Marker) (TYR/2024R), 1mg/mL, Clone: [TYR/2024R], Rabbit, Monoclonal BOT-BNUM2024-50

Artikelname	Tyrosinase (Melanoma Marker) (TYR/2024R), 1mg/mL, Clone: [TYR/2024R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2024-50
Hersteller Artikelnummer	BNUM2024-50
Alternativnummer	BOT-BNUM2024-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human tyrosinase (TYR) protein
Produktbeschreibung	This antibody recognizes a cluster of proteins between 70-80 kDa, identified as tyrosinase. Occasionally a minor band at 55 kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[TYR/2024R]
Molekulargewicht	70-80 kDa

UniProt	P14679
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 or 10 mM citrate buffer pH6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user