

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/1388), CF740 conjugate, 0.1mg/mL, IgG2a, Clone: [FTL/1388], CF 740, Mouse, Monoclonal BOT-BNC741388-100**

Artikelname	Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/1388), CF740 conjugate, 0.1mg/mL, IgG2a, Clone: [FTL/1388], CF 740, Mouse, Monoclonal
Artikelnummer	BOT-BNC741388-100
Hersteller Artikelnummer	BNC741388-100
Alternativnummer	BOT-BNC741388-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human FTL protein fragment (aa 38-165) (exact sequence is proprietary)
Konjugation	CF 740
Produktbeschreibung	Mammalian ferritins consist of 24 subunits made up of 2 types of polypeptide chains, ferritin heavy chain and ferritin light chain. Ferritin heavy chains catalyze the first step in iron storage, the oxidation of Fe (II), whereas ferritin light chains...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[FTL/1388]
Molekulargewicht	19-25 kDa
Isotyp	IgG2a
UniProt	<a href="#">P02792</a>
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 0.1-0.2 ug/mL Flow cytometry 0.1-0.2ug/million cells Immunofluorescence 0.1-0.2ug/ml Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user