

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

FSH-Receptor (Ovarian Marker) (FSHR/1400), CF740 conjugate, 0.1mg/mL, IgG1, Clone: [FSHR/1400], CF 740, Mouse, Monoclonal BOT-BNC741400-100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | FSH-Receptor (Ovarian Marker) (FSHR/1400), CF740 conjugate, 0.1mg/mL, IgG1, Clone: [FSHR/1400], CF 740, Mouse, Monoclonal |
| Artikelnummer | BOT-BNC741400-100 |
| Hersteller Artikelnummer | BNC741400-100 |
| Alternativnummer | BOT-BNC741400-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length FSHR protein |
| Konjugation | CF 740 |
| Produktbeschreibung | Follicle-stimulating hormone receptor (FSHR) is a 695 amino acid G protein coupled receptor. FSH binds to the receptor in a hand-clasp fashion via its alpha and beta subunits. While the beta subunit of FSH is involved in the binding of FSH to the rec... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [FSHR/1400] |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht | 75 kDa |
| Isotyp | IgG1 |
| UniProt | P23945 |
| Puffer | PBS, 0.1% rBSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |