

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (NX2/1422R), CF740 conjugate, 0.1mg/mL, Clone: [NX2/1422R], CF 740, Rabbit, Monoclonal BOT-BNC741422-500

Artikelname	NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (NX2/1422R), CF740 conjugate, 0.1mg/mL, Clone: [NX2/1422R], CF 740, Rabbit, Monoclonal
Artikelnummer	BOT-BNC741422-500
Hersteller Artikelnummer	BNC741422-500
Alternativnummer	BOT-BNC741422-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human NKX2.2 protein fragment (exact sequence is proprietary)
Konjugation	CF 740
Produktbeschreibung	Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[NX2/1422R]
Molekulargewicht	40-50 kDa
UniProt	O95096
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>