

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Wilms Tumor 1 (WT1) (Wilms Tumor & Mesothelial Marker) (WT1/1434R), CF740 conjugate, 0.1mg/mL, Clone: [WT1/1434R], CF 740, Rabbit, Monoclonal BOT-BNC741434-500

Artikelname	Wilms Tumor 1 (WT1) (Wilms Tumor & Mesothelial Marker) (WT1/1434R), CF740 conjugate, 0.1mg/mL, Clone: [WT1/1434R], CF 740, Rabbit, Monoclonal
Artikelnummer	BOT-BNC741434-500
Hersteller Artikelnummer	BNC741434-500
Alternativnummer	BOT-BNC741434-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length WT1 protein
Konjugation	CF 740
Produktbeschreibung	This antibody recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subse...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[WT1/1434R]
Molekulargewicht	47-55 kDa
UniProt	P19544
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user