

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Factor XIIIa (Coagulation Factor XIII A Chain) (F13A1/1447), CF740 conjugate, 0.1mg/mL, IgG2a, Clone: [F13A1/1447], CF 740, Mouse, Monoclonal BOT-BNC741447-500

| | |
|--------------------------|--|
| Artikelname | Factor XIIIa (Coagulation Factor XIII A Chain) (F13A1/1447), CF740 conjugate, 0.1mg/mL, IgG2a, Clone: [F13A1/1447], CF 740, Mouse, Monoclonal |
| Artikelnummer | BOT-BNC741447-500 |
| Hersteller Artikelnummer | BNC741447-500 |
| Alternativnummer | BOT-BNC741447-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is proprietary) |
| Konjugation | CF 740 |
| Produktbeschreibung | This antibody recognizes a protein of 83 kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macroph... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [F13A1/1447] |
| Molekulargewicht | 83 kDa |
| Isotyp | IgG2a |
| UniProt | P00488 |
| Puffer | PBS, 0.1% rBSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |