

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rat GADD45A protein, His tag, Unconjugated GTX00066-PRO

Artikelname	Rat GADD45A protein, His tag, Unconjugated
Artikelnummer	GTX00066-PRO
Hersteller Artikelnummer	GTX00066-pro
Alternativnummer	GTX00066-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Rat
Konjugation	Unconjugated
NCBI	25112
UniProt	P48317
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Met1~Arg165

Anwendungsbeschreibung

Growth arrest and DNA-damage-inducible protein GADD45 alpha (GADD45a) is a member of the GADD45 family. It may affect PCNA interaction with some CDK (cell division protein kinase) complexes, stimulates DNA excision repair in vitro and inhibits entry of cells into S phase. In T-cells, functions as a regulator of p38 MAPKs by inhibiting p38 phosphorylation and activity. Besides, Proliferating Cell Nuclear Antigen (PCNA) has been identified as an interactor of GADD45a, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat GADD45a and recombinant rat PCNA. Briefly, GADD45a were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to PCNA-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-GADD45a pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of GADD45a and PCNA was in a dose dependent manner.