

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human MCP4 protein, His tag (active), Unconjugated GTX00107-PRO

Artikelname	Human MCP4 protein, His tag (active), Unconjugated
Artikelnummer	GTX00107-PRO
Hersteller Artikelnummer	GTX00107-pro
Alternativnummer	GTX00107-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	6357
UniProt	Q99616
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Phe17~Thr98 (NP_005399.1)

Anwendungsbeschreibung	CCL13 (C-C motif chemokine 13) is a chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils, which belongs to the CC chemokine subfamily. It has been reported that CCL13 can induce chemotactic migration of THP-1 cells. Therefore, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of CCL13 on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (100 µl cell suspension, 1x10⁶ cells/ml in RPMI-1640 with 0.5% FBS) and CCL13 (50 ng/ml and 100 ng/ml diluted separately in serum free RPMI-1640) was added in lower chamber with a polycarbonate filter (8µm pore size) used to separate the two compartments. After incubation at 37°C with 5% CO₂ for 3h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (*100) and the number of migrated cells were counted at high magnification (*400) randomly. CCL13 is able to induce migration of THP-1 cells.
------------------------	---