

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse SOCS3 protein, His tag, Unconjugated GTX00304-PRO

Artikelname	Mouse SOCS3 protein, His tag, Unconjugated
Artikelnummer	GTX00304-PRO
Hersteller Artikelnummer	GTX00304-pro
Alternativnummer	GTX00304-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Mouse
Konjugation	Unconjugated
NCBI	12702
UniProt	O35718
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	Full length protein, N-terminal His-Tag, Met1~Leu225 (NP_031733.1)

Anwendungsbeschreibung	Suppressor of cytokine signaling 3 (SOCS3) is a member of the STAT-induced STAT inhibitor (SSI), also known as suppressor of cytokine signaling (SOCS), family that negatively regulates cytokine signal transduction. SOCS3 is feedback inhibitors of the Janus kinase (JAK) and signal transducer and activator of transcription (STAT) signaling pathway. Inhibits cytokine signal transduction by binding to tyrosine kinase receptors including gp130, LIF, erythropoietin, insulin, IL12, GCSF and leptin receptors. SOCS3 also can bind to JAK2 kinase inhibits its kinase activity and inhibits insulin signaling in adipose tissue and the liver. Besides, Glycoprotein 130 (gp130) has been identified as an interactor of SOCS3, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse SOCS3 and recombinant mouse gp130. Briefly, SOCS3 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to SOCS3-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-SOCS3 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of SOCS3 and gp130 was in a dose dependent manner.
------------------------	---