

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rat FGF13 protein, His tag (active), Unconjugated GTX00350-PRO

Artikelname	Rat FGF13 protein, His tag (active), Unconjugated
Artikelnummer	GTX00350-PRO
Hersteller Artikelnummer	GTX00350-pro
Alternativnummer	GTX00350-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Rat
Konjugation	Unconjugated
NCBI	84488
UniProt	Q9ERW3
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	Full length protein, N-terminal His-Tag, Met1~Thr192 (NP_445880.1)

Anwendungsbeschreibung

FGF13 (Fibroblast growth factor 13) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. A proliferation assay was conducted to detect the bioactivity of recombinant rat FGF13 using 3T3 cells. Briefly, 3T3 cells were seeded into triplicate wells of 96-well plates at a density of 2000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of FGF13. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of 3T3 cells after incubation with FGF13 for 48h observed by inverted microscope. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant FGF13 for 48h. And FGF13 significantly increased cell viability of 3T3 cells.