

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

FUT3 Antibody / Blood Group Lewis b, Clone: [SPM194], Mouse, Monoclonal NSJ-V2520-100UG

Artikelname	FUT3 Antibody / Blood Group Lewis b, Clone: [SPM194], Mouse, Monoclonal
Artikelnummer	NSJ-V2520-100UG
Hersteller Artikelnummer	V2520-100UG
Alternativnummer	NSJ-V2520-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Mucin isolated from a human ovarian cyst fluid was used as the immunogen for the FUT3 antibody.
Produktbeschreibung	The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metas...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM194]
UniProt	P21217

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the FUT3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.