

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

FUT3 Antibody / Blood Group Lewis b, Clone: [SPM194], Mouse, Monoclonal NSJ-V2520IHC-7ML

Artikelname	FUT3 Antibody / Blood Group Lewis b, Clone: [SPM194], Mouse, Monoclonal
Artikelnummer	NSJ-V2520IHC-7ML
Hersteller Artikelnummer	V2520IHC-7ML
Alternativnummer	NSJ-V2520IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Mucin isolated from a human ovarian cyst fluid was used as the immunogen for the FUT3 antibody.
Produktbeschreibung	The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metas...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM194]
UniProt	P21217

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the FUT3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.