

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD15 Antibody, Clone: [BRA-4F1], Mouse, Monoclonal NSJ-V2521SAF-100UG

| | |
|--------------------------|---|
| Artikelname | CD15 Antibody, Clone: [BRA-4F1], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2521SAF-100UG |
| Hersteller Artikelnummer | V2521SAF-100UG |
| Alternativnummer | NSJ-V2521SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Myelomonocytic leukemia cells were used as the immunogen for the CD15 antibody. |
| Produktbeschreibung | CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cell... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BRA-4F1] |
| UniProt | P22083 |
| Reinheit | PEG precipitation |

| | |
|------------------------|--|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 0.5-1ug/million cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the CD15 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes |