

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **SSEA-1 Antibody Cocktail, Clone: [FUT4/815 + BRA-4F1], Mouse, Monoclonal NSJ-V2524IHC-7ML**

Artikelname	SSEA-1 Antibody Cocktail, Clone: [FUT4/815 + BRA-4F1], Mouse, Monoclonal
Artikelnummer	NSJ-V2524IHC-7ML
Hersteller Artikelnummer	V2524IHC-7ML
Alternativnummer	NSJ-V2524IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Purified human neutrophils (FUT4/815) and Myelomonocytic leukemia cells (BRA-4F1) were used as the immunogen for the SSEA-1 antibody cocktail.
Produktbeschreibung	SSEA-1/CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, it is expressed in Reed-Sternberg...
Klonalität	Monoclonal
Klon-Bezeichnung	[FUT4/815 + BRA-4F1]
UniProt	<a href="#">P22083</a>

Reinheit	PEG precipitation
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the SSEA-1 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.