

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD57 Antibody, Clone: [SPM129], Mouse, Monoclonal NSJ-V2533-20UG

Artikelname	CD57 Antibody, Clone: [SPM129], Mouse, Monoclonal
Artikelnummer	NSJ-V2533-20UG
Hersteller Artikelnummer	V2533-20UG
Alternativnummer	NSJ-V2533-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human peripheral blood mononuclear cells were used as the immunogen for the CD57 antibody.
Produktbeschreibung	Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including c...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM129]
UniProt	Q9P2W7
Reinheit	PEG precipitation

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the CD57 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.