

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glypican-3 Antibody, Clone: [GPC3/863], Mouse, Monoclonal NSJ-V2538-20UG

Artikelname	Glypican-3 Antibody, Clone: [GPC3/863], Mouse, Monoclonal
Artikelnummer	NSJ-V2538-20UG
Hersteller Artikelnummer	V2538-20UG
Alternativnummer	NSJ-V2538-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant full-length human GPC3 protein was used as the immunogen for the Glypican-3 antibody.
Produktbeschreibung	Glypican-3 (GPC3) is a glycosylphosphatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, mel...
Klonalität	Monoclonal
Klon-Bezeichnung	[GPC3/863]
UniProt	P51654
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/million cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>Optimal dilution of the Glypican-3 antibody should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. 3. View the recombinant version of this recombinant-gpc3-antibody-rgpc3-863-v3456 Glypican-3 antibody.