

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### MFGE8 Antibody, Clone: [MFG-06], Mouse, Monoclonal NSJ-V2703IHC-7ML

Artikelname	MFGE8 Antibody, Clone: [MFG-06], Mouse, Monoclonal
Artikelnummer	NSJ-V2703IHC-7ML
Hersteller Artikelnummer	V2703IHC-7ML
Alternativnummer	NSJ-V2703IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	A human milk fat globule membrane preparation was used as the immunogen for the MFGE8 antibody.
Produktbeschreibung	Recognizes a protein of 40-45kDa, identified as human milk fat globule membrane protein (HMFG), Lactadherin, or MFG-E8. HMFG is present on normal human breast epithelial cells and cell lines derived from breast carcinomas, as well as to the outer sur...
Klonalität	Monoclonal
Klon-Bezeichnung	[MFG-06]
UniProt	<a href="#">Q08431</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the MFGE8 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.