

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MFG-E8 Antibody, Clone: [SPM291], Mouse, Monoclonal NSJ-V2704SAF-100UG

| | |
|--------------------------|---|
| Artikelname | MFG-E8 Antibody, Clone: [SPM291], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2704SAF-100UG |
| Hersteller Artikelnummer | V2704SAF-100UG |
| Alternativnummer | NSJ-V2704SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | A human milk fat globule membrane preparation was used as the immunogen for the MFG-E8 antibody. |
| Produktbeschreibung | Recognizes a protein of 40-45kDa, identified as human milk fat globule membrane protein (HMFG), Lactadherin or MFG-E8. HMFG is present on normal human breast epithelial cells and cell lines derived from breast carcinomas, as well as to the outer surf... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM291] |
| UniProt | Q08431 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the MFG-E8 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |