

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Lactadherin Antibody, Clone: [EDM45], Mouse, Monoclonal NSJ-V2705SAF-100UG

Artikelname	Lactadherin Antibody, Clone: [EDM45], Mouse, Monoclonal
Artikelnummer	NSJ-V2705SAF-100UG
Hersteller Artikelnummer	V2705SAF-100UG
Alternativnummer	NSJ-V2705SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	A delipidated human milk fat globule membrane preparation was used as the immunogen for the Lactadherin antibody.
Produktbeschreibung	Recognizes a protein of 40-45kDa, identified as human milk fat globule membrane protein (HMFG), Lactadherin or MFG-EA. HMFG is present on normal human breast epithelial cells and cell lines derived from breast carcinomas, as well as to the outer surf...
Klonalität	Monoclonal
Klon-Bezeichnung	[EDM45]
UniProt	Q08431
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Lactadherin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.