

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD99 Antibody, Clone: [SPM586], Mouse, Monoclonal NSJ-V2707IHC-7ML

Artikelname	CD99 Antibody, Clone: [SPM586], Mouse, Monoclonal
Artikelnummer	NSJ-V2707IHC-7ML
Hersteller Artikelnummer	V2707IHC-7ML
Alternativnummer	NSJ-V2707IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Purified E-rosette forming cells from human peripheral blood lymphocytes were used as the immunogen for the CD99 antibody.
Produktbeschreibung	Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. This antigen is expressed on the cell membrane of some lymphocytes, cortical thymocytes, and granulosa cells of the ovary. Most pancreatic islet cell...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM586]
UniProt	<a href="#">P14209</a>
Reinheit	PEG precipitation

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the CD99 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.