

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD99 Antibody, Clone: [MIC2/877], Mouse, Monoclonal NSJ-V2709SAF-100UG

Artikelname	CD99 Antibody, Clone: [MIC2/877], Mouse, Monoclonal
Artikelnummer	NSJ-V2709SAF-100UG
Hersteller Artikelnummer	V2709SAF-100UG
Alternativnummer	NSJ-V2709SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the CD99 antibody.
Produktbeschreibung	Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by a...
Klonalität	Monoclonal
Klon-Bezeichnung	[MIC2/877]
UniProt	<a href="#">P14209</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at Room Temp
Anwendungsbeschreibung	Optimal dilution of the CD99 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.