

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 Antibody, Clone: [139H2], Mouse, Monoclonal NSJ-V2726IHC-7ML

Artikelname	MUC1 Antibody, Clone: [139H2], Mouse, Monoclonal
Artikelnummer	NSJ-V2726IHC-7ML
Hersteller Artikelnummer	V2726IHC-7ML
Alternativnummer	NSJ-V2726IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human milk-fat globule membranes (HMFGM) was used as the immunogen for the MUC1 antibody.
Produktbeschreibung	This mAb reacts with MUC1/Mucin-1/Epithelial Marker Antigen/EMA. The dominant epitope of this mAb has not yet been determined. MUC1 is a large cell surface mucin glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoie...
Klonalität	Monoclonal
Klon-Bezeichnung	[139H2]
UniProt	P15941
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the MUC1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.