

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-MUC-1 Antibody, Clone: [SPM492], Mouse, Monoclonal NSJ-V2727SAF-100UG

Artikelname	Anti-MUC-1 Antibody, Clone: [SPM492], Mouse, Monoclonal
Artikelnummer	NSJ-V2727SAF-100UG
Hersteller Artikelnummer	V2727SAF-100UG
Alternativnummer	NSJ-V2727SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Breast cancer cell line ZR75-1 was used as the immunogen for the anti-MUC-1 antibody.
Produktbeschreibung	This mAb reacts with MUC1/Mucin-1/Epithelial Marker Antigen/EMA. MUC1 is a large cell surface mucin glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoietic cell lineages. It is expressed on most secretory epitheliu...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM492]
UniProt	P15941
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>Optimal dilution of the anti-MUC-1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>