

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin-3 Antibody, Clone: [M3.1], Mouse, Monoclonal NSJ-V2730IHC-7ML

Artikelname	Mucin-3 Antibody, Clone: [M3.1], Mouse, Monoclonal
Artikelnummer	NSJ-V2730IHC-7ML
Hersteller Artikelnummer	V2730IHC-7ML
Alternativnummer	NSJ-V2730IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids (C-HSTPSFTSSITTTETTSHTPSFTSSITTTETTS) which contains two of the MUC3 tandem repeats was used as the immunogen for the Mucin-3 antibody.
Produktbeschreibung	It recognizes a protein of HMW, identified as mucin 3 glycoprotein (MUC3). Its epitope localizes between amino acids SITTE. This mAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC2. MUC3 is distributed in colon and rectu...
Klonalität	Monoclonal
Klon-Bezeichnung	[M3.1]
UniProt	Q02505
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Mucin-3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.