

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-MUC3 Antibody, Clone: [SPM200], Mouse, Monoclonal NSJ-V2731IHC-7ML

Artikelname	Anti-MUC3 Antibody, Clone: [SPM200], Mouse, Monoclonal
Artikelnummer	NSJ-V2731IHC-7ML
Hersteller Artikelnummer	V2731IHC-7ML
Alternativnummer	NSJ-V2731IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids (C-HSTPSFTSSITTTETTSHTPSFTSSITTTETTS) which contains two of the MUC3 tandem repeats was used as the immunogen for the anti-MUC3 antibody.
Produktbeschreibung	MUC3 is distributed in colon and rectum, and is also present to a lesser extent in breast, lung and salivary gland tissues. The Mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of ta...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM200]
UniProt	<a href="#">Q02505</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the anti-MUC3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.