

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC5-AC Antibody, Clone: [SPM488], Mouse, Monoclonal NSJ-V2737IHC-7ML

Artikelname	MUC5-AC Antibody, Clone: [SPM488], Mouse, Monoclonal
Artikelnummer	NSJ-V2737IHC-7ML
Hersteller Artikelnummer	V2737IHC-7ML
Alternativnummer	NSJ-V2737IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	A synthetic peptide of human MUC5AC tandem repeat was used as the immunogen for the MUC5-AC antibody.
Produktbeschreibung	Mucin 5AC glycoprotein (MUC5AC) is a HMW glycoprotein belonging to the superfamily of mucins. Mucins are produced by epithelial cells and can be divided into two families, secretory mucins and membrane bound mucins. MUC5AC is a mucus-forming, secrete...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM488]
UniProt	P98088
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the MUC5-AC antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.