

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MUC5AC Antibody, Clone: [MUC5AC/917], Mouse, Monoclonal NSJ-V2738-20UG**

Artikelname	MUC5AC Antibody, Clone: [MUC5AC/917], Mouse, Monoclonal
Artikelnummer	NSJ-V2738-20UG
Hersteller Artikelnummer	V2738-20UG
Alternativnummer	NSJ-V2738-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human partial protein was used as the immunogen for the MUC5AC antibody.
Produktbeschreibung	Mucin 5AC glycoprotein (MUC5AC) is a 641kDa glycoprotein belonging to the superfamily of mucins. Mucins are high molecular weight glycoproteins produced by epithelial cells and can be divided into two families, secretory mucins and membrane bound muc...
Klonalität	Monoclonal
Klon-Bezeichnung	[MUC5AC/917]
UniProt	<a href="#">P98088</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)
Anwendungsbeschreibung	Optimal dilution of the MUC5AC antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.