

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MUC5AC Antibody Cocktail, Clone: [MUC5AC/917 + 45M1], Mouse, Monoclonal NSJ-V2740SAF-100UG**

Artikelname	MUC5AC Antibody Cocktail, Clone: [MUC5AC/917 + 45M1], Mouse, Monoclonal
Artikelnummer	NSJ-V2740SAF-100UG
Hersteller Artikelnummer	V2740SAF-100UG
Alternativnummer	NSJ-V2740SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein (MUC5AC/917) and an M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient (45M1) were used as the immunogen for the MUC5AC antibody cocktail.
Produktbeschreibung	Mucin 5AC glycoprotein (MUC5AC) is a 641kDa glycoprotein belonging to the superfamily of mucins. Mucins are high molecular weight glycoproteins produced by epithelial cells and can be divided into two families, secretory mucins and membrane bound muc...
Klonalität	Monoclonal
Klon-Bezeichnung	[MUC5AC/917 + 45M1]

UniProt	<a href="#">P98088</a>
Reinheit	Protein G affinity chromatography
Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>Optimal dilution of the MUC5AC antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>