

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Antibody, Clone: [9E10.3], Mouse, Monoclonal NSJ-V2744IHC-7ML

Artikelname	c-Myc Antibody, Clone: [9E10.3], Mouse, Monoclonal
Artikelnummer	NSJ-V2744IHC-7ML
Hersteller Artikelnummer	V2744IHC-7ML
Alternativnummer	NSJ-V2744IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA were used as the immunogen for the c-Myc antibody.
Produktbeschreibung	It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This mAb shows no cross-reaction with v-myc. c-myc...
Klonalität	Monoclonal
Klon-Bezeichnung	[9E10.3]
UniProt	P01106
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the c-Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.