

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Antibody, Clone: [9E10.3], Mouse, Monoclonal NSJ-V2744SAF-100UG

Artikelname	c-Myc Antibody, Clone: [9E10.3], Mouse, Monoclonal
Artikelnummer	NSJ-V2744SAF-100UG
Hersteller Artikelnummer	V2744SAF-100UG
Alternativnummer	NSJ-V2744SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA were used as the immunogen for the c-Myc antibody.
Produktbeschreibung	It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This mAb shows no cross-reaction with v-myc. c-myc...
Klonalität	Monoclonal
Klon-Bezeichnung	[9E10.3]
UniProt	P01106
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the c-Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.