

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myc Antibody, Clone: [SPM237], Mouse, Monoclonal NSJ-V2745-20UG

| | |
|--------------------------|--|
| Artikelname | Myc Antibody, Clone: [SPM237], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2745-20UG |
| Hersteller Artikelnummer | V2745-20UG |
| Alternativnummer | NSJ-V2745-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA from the C-terminus were used as the immunogen for the Myc antibody. |
| Produktbeschreibung | Myc, or c-Myc, is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-Myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM237] |
| UniProt | P01106 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2) |
| Anwendungsbeschreibung | Optimal dilution of the Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |