

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Myc Antibody, Clone: [SPM237], Mouse, Monoclonal NSJ-V2745SAF-100UG**

|                          |  |
|--------------------------|--|
| Artikelname              | Myc Antibody, Clone: [SPM237], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V2745SAF-100UG   |
| Hersteller Artikelnummer | V2745SAF-100UG   |
| Alternativnummer         | NSJ-V2745SAF-100UG   |
| Hersteller               | NSJ Bioreagents  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | IHC-P  |
| Spezies Reaktivität      | Human  |
| Immunogen                | Amino acids AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA from the C-terminus were used as the immunogen for the Myc antibody.  |
| Produktbeschreibung      | Myc, or c-Myc, is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-Myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia... |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [SPM237]   |
| UniProt                  | <a href="#">P01106</a>   |
| Reinheit                 | Protein G affinity chromatography  |

|                        |   |
|------------------------|---|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free  |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)  |
| Anwendungsbeschreibung | Optimal dilution of the Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |