

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Antibody Cocktail, Clone: [MYC275 + MYC909], Mouse, Monoclonal NSJ-V2747-100UG

Artikelname	c-Myc Antibody Cocktail, Clone: [MYC275 + MYC909], Mouse, Monoclonal
Artikelnummer	NSJ-V2747-100UG
Hersteller Artikelnummer	V2747-100UG
Alternativnummer	NSJ-V2747-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the c-Myc antibody cocktail.
Produktbeschreibung	It recognizes a transcription factor of 64-67kDa, identified as c-myc. This mAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a variety of tumo...
Klonalität	Monoclonal
Klon-Bezeichnung	[MYC275 + MYC909]
UniProt	P01106
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the c-Myc antibody cocktail should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min