

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

SM-MHC Antibody, Clone: [SPM201], Mouse, Monoclonal NSJ-V2749SAF-100UG

| | |
|--------------------------|---|
| Artikelname | SM-MHC Antibody, Clone: [SPM201], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2749SAF-100UG |
| Hersteller Artikelnummer | V2749SAF-100UG |
| Alternativnummer | NSJ-V2749SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Human uterus extract was used as the immunogen for the SM-MHC antibody. |
| Produktbeschreibung | Smooth muscle myosin heavy chain (SM-MHC) is a cytoplasmic structural protein, which is a major component of the contractile apparatus in smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated, appearing early in smooth ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM201] |
| UniProt | P35749 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | <p>The stated application concentrations are suggested starting amounts. Optimal dilution of the SM-MHC antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |