

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

SM-MHC Antibody, Clone: [ID8], Mouse, Monoclonal NSJ-V2750-20UG

| | |
|--------------------------|---|
| Artikelname | SM-MHC Antibody, Clone: [ID8], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2750-20UG |
| Hersteller Artikelnummer | V2750-20UG |
| Alternativnummer | NSJ-V2750-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Human uterus extract was used as the immunogen for the SM-MHC antibody. |
| Produktbeschreibung | SM-MHC, or Myosin-11, is a smooth muscle myosin belonging to the myosin heavy chain family. Myosin-11 is a subunit of a hexameric protein that consists of two heavy chain subunits and two pairs of non-identical light chain subunits. It is a major con... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [ID8] |
| UniProt | P35749 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the SM-MHC antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |