

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### SM-MHC Antibody, Clone: [ID8], Mouse, Monoclonal NSJ-V2750IHC-7ML

Artikelname	SM-MHC Antibody, Clone: [ID8], Mouse, Monoclonal
Artikelnummer	NSJ-V2750IHC-7ML
Hersteller Artikelnummer	V2750IHC-7ML
Alternativnummer	NSJ-V2750IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Human uterus extract was used as the immunogen for the SM-MHC antibody.
Produktbeschreibung	SM-MHC, or Myosin-11, is a smooth muscle myosin belonging to the myosin heavy chain family. Myosin-11 is a subunit of a hexameric protein that consists of two heavy chain subunits and two pairs of non-identical light chain subunits. It is a major con...
Klonalität	Monoclonal
Klon-Bezeichnung	[ID8]
UniProt	<a href="#">P35749</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the SM-MHC antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.