

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NCAM Antibody, Clone: [NCAM1/784], Mouse, Monoclonal NSJ-V2753IHC-7ML

| | |
|--------------------------|---|
| Artikelname | NCAM Antibody, Clone: [NCAM1/784], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2753IHC-7ML |
| Hersteller Artikelnummer | V2753IHC-7ML |
| Alternativnummer | NSJ-V2753IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human protein was used as the immunogen for the NCAM antibody. |
| Produktbeschreibung | This mAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [NCAM1/784] |
| UniProt | P13591 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the NCAM antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |