

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **NCL Antibody / Nucleolin, Clone: [364-5], Mouse, Monoclonal NSJ-V2757SAF-100UG**

Artikelname	NCL Antibody / Nucleolin, Clone: [364-5], Mouse, Monoclonal
Artikelnummer	NSJ-V2757SAF-100UG
Hersteller Artikelnummer	V2757SAF-100UG
Alternativnummer	NSJ-V2757SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	SUDHL1 cell nuclear lysate was used as the immunogen for the NCL antibody.
Produktbeschreibung	Recognizes a protein of ~76kDa, which is identified as Nucleolin (NCL). It is the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleola...
Klonalität	Monoclonal
Klon-Bezeichnung	[364-5]
UniProt	<a href="#">P19338</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 0.5-1ug/ml, Western blot: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the NCL antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.