

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Nucleolin Antibody Cocktail, Clone: [364-5 + NCL/902], Mouse, Monoclonal NSJ-V2758SAF-100UG

| | |
|--------------------------|---|
| Artikelname | Nucleolin Antibody Cocktail, Clone: [364-5 + NCL/902], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2758SAF-100UG |
| Hersteller Artikelnummer | V2758SAF-100UG |
| Alternativnummer | NSJ-V2758SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | SUDHL1 cell nuclear lysate (364-5) and recombinant human protein (NCL/902) were used as the immunogen for the Nucleolin antibody cocktail. |
| Produktbeschreibung | Recognizes a protein of ~76kDa, which is identified as Nucleolin (NCL). It is the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleola... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [364-5 + NCL/902] |
| UniProt | P19338 |

| | |
|------------------------|---|
| Reinheit | Protein G affinity chromatography |
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Western blot: 0.5-1ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the Nucleolin antibody cocktail should be determined by the researcher. 1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |