

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NF-H Antibody (phospho), Clone: [NE14], Mouse, Monoclonal NSJ-V2759IHC-7ML

Artikelname	NF-H Antibody (phospho), Clone: [NE14], Mouse, Monoclonal
Artikelnummer	NSJ-V2759IHC-7ML
Hersteller Artikelnummer	V2759IHC-7ML
Alternativnummer	NSJ-V2759IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Crude neurofilament preparation from porcine spinal cord was used as the immunogen for the NF-H antibody (phospho).
Produktbeschreibung	This mAb reacts with a 200kDa protein, identified as heavy sub-unit of neurofilaments (NF-H). It reacts specifically with the phosphorylated KSP/KEP segment at the C-terminus of the heavy subunit (NF-H) of neurofilaments. After dephosphorylation of n...
Klonalität	Monoclonal
Klon-Bezeichnung	[NE14]
UniProt	P12036
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the NF-H antibody (phospho) should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.