

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NGF Receptor Antibody, Clone: [SPM299], Mouse, Monoclonal NSJ-V2764IHC-7ML

Artikelname	NGF Receptor Antibody, Clone: [SPM299], Mouse, Monoclonal
Artikelnummer	NSJ-V2764IHC-7ML
Hersteller Artikelnummer	V2764IHC-7ML
Alternativnummer	NSJ-V2764IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	NGFR from A875 melanoma cells was used as the immunogen for the NGF Receptor antibody. Its epitope is within amino acids 1-160 of the extracellular domain of NGFR/NTR.
Produktbeschreibung	It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGF-receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine resi...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM299]
UniProt	P08138
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the NGF Receptor antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.