

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **NGF Receptor Antibody Cocktail, Clone: [NGFR5 + NTR/912], Mouse, Monoclonal NSJ-V2766IHC-7ML**

Artikelname	NGF Receptor Antibody Cocktail, Clone: [NGFR5 + NTR/912], Mouse, Monoclonal
Artikelnummer	NSJ-V2766IHC-7ML
Hersteller Artikelnummer	V2766IHC-7ML
Alternativnummer	NSJ-V2766IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	NGFR from A875 melanoma cells (NGFR5) and recombinant human protein (NTR/912) were used as the immunogen for the NGF Receptor antibody cocktail.
Produktbeschreibung	It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neurobl...
Klonalität	Monoclonal
Klon-Bezeichnung	[NGFR5 + NTR/912]
UniProt	<a href="#">P08138</a>

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the NGF Receptor antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.