

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ornithine Decarboxylase Antibody, Clone: [ODC1/487], Mouse, Monoclonal NSJ-V2770SAF-100UG

Artikelname	Ornithine Decarboxylase Antibody, Clone: [ODC1/487], Mouse, Monoclonal
Artikelnummer	NSJ-V2770SAF-100UG
Hersteller Artikelnummer	V2770SAF-100UG
Alternativnummer	NSJ-V2770SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human protein was used as the immunogen for the Ornithine Decarboxylase antibody.
Produktbeschreibung	Recognizes a 53kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activity...
Klonalität	Monoclonal
Klon-Bezeichnung	[ODC1/487]
UniProt	P11926
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml, Western blot: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Ornithine Decarboxylase antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.