

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **PAX6 Antibody, Clone: [PAX6/1166], Mouse, Monoclonal NSJ-V2771IHC-7ML**

Artikelname	PAX6 Antibody, Clone: [PAX6/1166], Mouse, Monoclonal
Artikelnummer	NSJ-V2771IHC-7ML
Hersteller Artikelnummer	V2771IHC-7ML
Alternativnummer	NSJ-V2771IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	A recombinant fragment (N-terminus, aa 1-300) was used as the immunogen for the PAX6 antibody.
Produktbeschreibung	Pax genes contain paired domains with strong homology to genes in Drosophila, which are involved in programming early development. Lesions in the Pax-6 gene account for most cases of aniridia, a congenital malformation of the eye, chiefly characteriz...
Klonalität	Monoclonal
Klon-Bezeichnung	[PAX6/1166]
UniProt	<a href="#">P26367</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the PAX6 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.