

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### PD-1 Antibody, Clone: [PDCD1/922], Mouse, Monoclonal NSJ-V2776-20UG

Artikelname	PD-1 Antibody, Clone: [PDCD1/922], Mouse, Monoclonal
Artikelnummer	NSJ-V2776-20UG
Hersteller Artikelnummer	V2776-20UG
Alternativnummer	NSJ-V2776-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the PD-1 antibody.
Produktbeschreibung	PDCD-1 (programmed cell death-1 protein), also called PD-1 and CD279, is a type I transmembrane receptor and a member of the immunoglobulin gene superfamily. It is expressed on activated T-cells, B-cells, and myeloid cells. Anti-PDCD-1 is a marker of a...
Klonalität	Monoclonal
Klon-Bezeichnung	[PDCD1/922]
UniProt	<a href="#">Q15116</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the PD-1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.