

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PECAM Antibody, Clone: [1A10], Mouse, Monoclonal NSJ-V2781IHC-7ML

Artikelname	PECAM Antibody, Clone: [1A10], Mouse, Monoclonal
Artikelnummer	NSJ-V2781IHC-7ML
Hersteller Artikelnummer	V2781IHC-7ML
Alternativnummer	NSJ-V2781IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Prokaryotic recombinant fusion protein corresponding to a portion of the extracellular domain downstream of the signal sequence of CD31/PECAM was used as the immunogen for the PECAM antibody.
Produktbeschreibung	CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime...
Klonalität	Monoclonal
Klon-Bezeichnung	[1A10]
UniProt	P16284
Reinheit	Unpurified high titer supernatant

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the PECAM antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.