

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytochrome C Antibody, Clone: [7H8.2C12], Mouse, Monoclonal NSJ-V2786-20UG

| | |
|--------------------------|---|
| Artikelname | Cytochrome C Antibody, Clone: [7H8.2C12], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2786-20UG |
| Hersteller Artikelnummer | V2786-20UG |
| Alternativnummer | NSJ-V2786-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Synthetic peptides corresponding to amino acid 1-80, 81-104 and 66-104 of pigeon Cytochrome C was used as the immunogen for the Cytochrome C antibody. |
| Produktbeschreibung | It recognizes an epitope within amino acids 93-104 of pigeon Cytochrome C, a well-characterized mobile electron transport protein that is essential to energy conversion in all aerobic organisms. In mammalian cells, this highly conserved protein is no... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [7H8.2C12] |
| UniProt | P99999 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the Cytochrome C antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |